Warfarin (Coumadin) Dosing Guide

FOR 6 mg TABLETS

Intermountain® Healthcare

Tablet Strength: 6 milligrams (mg) Target PT/INR: _____

Warfarin (brand name **Coumadin**) is prescribed in a weekly dose. Coumadin tablets come in multiple strengths. You are on a **6 mg tablet**. The table below shows you how much to take each day of the week to add up to the correct weekly dose. Some days you may need to take ½ a tablet. ***It's important for you to take the exact dose you are prescribed, at the same time everyday — in the evening.***

Dosing for 6 mg warfarin (Coumadin) tablet							
Weekly Dose	Tablets per day (½ = a half tablet; 1 = 1 tablet; 1½ = 1½ tablets)						
	Sun	Mon	Tues	Wed	Thurs	Fri	Sat
☐ 21 mg	½	½	½	½	½	½	½
☐ 24 mg	1	½	½	½	½	½	½
☐ 27 mg	1	½	½	½	1	½	½
☐ 30 mg	1	½	1	½	1	½	½
☐ 33 mg	1	½	1	½	1	½	1
☐ 36 mg	1	½	1	1	1	½	1
☐ 39 mg	1	½	1	1	1	1	1
☐ 42 mg	1	1	1	1	1	1	1
☐ 45 mg	1½	1	1	1	1	1	1
☐ 48 mg	1½	1	1	1	1½	1	1
☐ 51 mg	1½	1	1½	1	1½	1	1
☐ 54 mg	1½	1	1½	1	1½	1	1½
☐ 57 mg	1½	1	1½	1½	1½	1	1½
☐ 60 mg	1½	1	1½	1½	1½	1½	1½
☐ 63 mg	1½	1½	1½	1½	1½	1½	1½
☐ 66 mg	2	1½	1½	1½	1½	1½	1½
☐ 69 mg	2	1½	1½	1½	2	1½	1½
☐ 71 mg	2	1½	2	1½	2	1½	1½
☐ 74 mg	2	1½	2	1½	2	1½	2
☐ 77 mg	2	1½	2	2	2	1 ½	2
☐ 80 mg	2	1 ½	2	2	2	2	2
☐ 84 mg	2	2	2	2	2	2	2

How to use this chart: Find the number in the left-most column that corresponds to your total weekly dose of Coumadin. Move across that row to see the individual dose for each day of the week. *(For example, if you are using the **6 mg** tablet and your total weekly dose is **27 mg**, you would take 1 whole tablet Sunday and Thursday, and ½ tablet the other days of the week.)*

The above information is not a substitute for professional medical advice. Please talk with your healthcare provider if you have any questions or concerns.

©2006-2011 Intermountain Healthcare. All rights reserved. Patient and Provider Publications (801) 442-2963 CAC060 - 08/11

Reminders about Coumadin and other medicines

- **Tell ALL care providers ALL medicines you are on.** Many medicines affect how Coumadin works. Be sure to tell our clinic staff ALL medicines you are taking — and be sure to tell your other care providers that you are taking Coumadin.

- **Call the clinic if you are prescribed any of the following medicines; you may need an adjustment to your Coumadin dose:**
  - Amiodarone (Pacerone or Cordarone)
  - Fluconazole (Diflucan)
  - Any antibiotic for more than a 3-day course

- **Get a PT/INR test in 7 days if you start, stop, or change dosage of any of the following medicines:**
  - Any antidepressant (medicine for depression)
  - Any statin (for example, Lipitor, Zocor, Pravachol, and Vytorin)
  - Any steroid (for example, Prednisone)
  - Interferon

- **Avoid aspirin and other non-steroidal anti-inflammatory drugs** such as ibuprofen unless prescribed by your physician. Tylenol (acetaminophen) is okay up to 1,000 mg (3 regular or 2 extra-strength) daily. Doses higher than that may alter the affect of Coumadin.

  Note: Call the clinic if you increase your regular dose of more than 1,000 mg/day of acetaminophen (Tylenol) or if you increase your dose of any other analgesic.

- **Don't take herbal supplements.** Many herbal medications affect how well Coumadin works. Never take an herbal supplement without talking with your doctor first.

Intermountain℠ Healthcare

The above information is not a substitute for professional medical advice. Please talk with your healthcare provider if you have any questions or concerns.

©2006-2011 Intermountain Healthcare. All rights reserved. Patient and Provider Publications (801) 442-2963 CAC060 - 08/11