

CORRECTION DOSE

This is the insulin dose you take for high blood glucose at meals and bedtime.



Meal time: 1 unit per 100mg/dL over 200

Bedtime: 1 unit per 100mg/dL over 300

Blood glucose	Before a meal	Bedtime
201–300	1	0
301–400	2	1
401–499	3	2
over 499	4	3

Only give correction dose if it has been **more than 3 hours** since the last dose of fast-acting insulin.

CORRECTION DOSE

This is the insulin dose you take for high blood glucose at meals and bedtime.



Meal time: 1 unit per 100mg/dL over 200

Bedtime: 1 unit per 100mg/dL over 300

Blood glucose	Before a meal	Bedtime
201–300	1	0
301–400	2	1
401–499	3	2
over 499	4	3

Only give correction dose if it has been **more than 3 hours** since the last dose of fast-acting insulin.

CORRECTION DOSE

This is the insulin dose you take for high blood glucose at meals and bedtime.



Meal time: 1 unit per 100mg/dL over 200

Bedtime: 1 unit per 100mg/dL over 300

Blood glucose	Before a meal	Bedtime
201–300	1	0
301–400	2	1
401–499	3	2
over 499	4	3

Only give correction dose if it has been **more than 3 hours** since the last dose of fast-acting insulin.

CORRECTION DOSE

This is the insulin dose you take for high blood glucose at meals and bedtime.



Meal time: 1 unit per 100mg/dL over 200

Bedtime: 1 unit per 100mg/dL over 300

Blood glucose	Before a meal	Bedtime
201–300	1	0
301–400	2	1
401–499	3	2
over 499	4	3

Only give correction dose if it has been **more than 3 hours** since the last dose of fast-acting insulin.

CORRECTION DOSE

This is the insulin dose you take for high blood glucose at meals and bedtime.



Meal time: 1 unit per 100mg/dL over 200

Bedtime: 1 unit per 100mg/dL over 300

Blood glucose	Before a meal	Bedtime
201–300	1	0
301–400	2	1
401–499	3	2
over 499	4	3

Only give correction dose if it has been **more than 3 hours** since the last dose of fast-acting insulin.

CORRECTION DOSE

This is the insulin dose you take for high blood glucose at meals and bedtime.



Meal time: 1 unit per 100mg/dL over 200

Bedtime: 1 unit per 100mg/dL over 300

Blood glucose	Before a meal	Bedtime
201–300	1	0
301–400	2	1
401–499	3	2
over 499	4	3

Only give correction dose if it has been **more than 3 hours** since the last dose of fast-acting insulin.



**Intermountain
Primary Children's Hospital**
The Child First and Always®

intermountainhealthcare.org/diabetes

primarychildrens.org/diabetes

© 2017 Intermountain Healthcare Inc.
All rights reserved
Primary Children's Hospital
PESDMC035_E



**Intermountain
Primary Children's Hospital**
The Child First and Always®

intermountainhealthcare.org/diabetes

primarychildrens.org/diabetes

© 2017 Intermountain Healthcare Inc.
All rights reserved
Primary Children's Hospital
PESDMC035_E



**Intermountain
Primary Children's Hospital**
The Child First and Always®

intermountainhealthcare.org/diabetes

primarychildrens.org/diabetes

© 2017 Intermountain Healthcare Inc.
All rights reserved
Primary Children's Hospital
PESDMC035_E



**Intermountain
Primary Children's Hospital**
The Child First and Always®

intermountainhealthcare.org/diabetes

primarychildrens.org/diabetes

© 2017 Intermountain Healthcare Inc.
All rights reserved
Primary Children's Hospital
PESDMC035_E



**Intermountain
Primary Children's Hospital**
The Child First and Always®

intermountainhealthcare.org/diabetes

primarychildrens.org/diabetes

© 2017 Intermountain Healthcare Inc.
All rights reserved
Primary Children's Hospital
PESDMC035_E



**Intermountain
Primary Children's Hospital**
The Child First and Always®

intermountainhealthcare.org/diabetes

primarychildrens.org/diabetes

© 2017 Intermountain Healthcare Inc.
All rights reserved
Primary Children's Hospital
PESDMC035_E