

Maternal Emotional Wellness After Loss:

Self-assessment and referral guide

A pregnancy loss can make it difficult for you to take care of your needs and emotional wellness. The information below can help you determine your level of emotional wellness and symptoms of concern, and to know Intermountain Healthcare's recommended actions. See <u>page 2</u> for information about community resources that are available to help you.

Low-risk symptoms	Recommended action
 □ Some sadness or low mood □ Some worry or feelings of anxiety □ Periodic crying or fluctuating mood □ More sleep struggles than expected □ Some disinterest in social activities 	 Give yourself permission to talk about your feelings. Don't be afraid to ask for or accept help. Get plenty of uninterrupted sleep. Eat nutritious foods every 3 to 4 hours and stay hydrated. Avoid caffeine and sweets when possible. Continue taking a prenatal vitamin for its emotional and physical benefits. Begin mild to moderate exercise as your doctor permits. Find ways to laugh and spend time with other adults.
Moderate-risk symptoms	Recommended action
 □ Excessive or uncontrollable crying □ Significant anxiety or panic attacks (symptoms may include rapid breathing and excessive sweating) □ Intense irritability and anger □ Intense feelings of shame or inadequacy □ Significant change in appetite or weight □ Difficulty concentrating or making decisions □ Less interest in favorite activities □ Concerns expressed by people close to you □ Moderate disturbances to appetite or sleep 	 Contact your obstetric (OB) provider and schedule an appointment as soon as possible. Seek psychiatric care as directed by your OB provider. Find a qualified provider or therapist by calling United Way's Help Me Grow at 801-691-5322 or contact your insurance provider for help accessing mental health services. Seek in-person or online support groups through social media. Educate yourself about postpartum mental health; postpartum.net is a great resource. Prioritize your needs in the home.
High-risk symptoms	Recommended action
 □ Strange beliefs or visual disturbances □ Thoughts to harm yourself □ Taking actions to harm yourself □ Obsessive, intrusive thoughts about your pregnancy □ Significant harmful reactions to medication □ Confusion □ Neglect 	 Consult with your OB provider immediately. Go to an emergency room right away if you fear for your own safety. Remember you are not alone and help is available. Speak openly and honestly about your symptoms. Tell loved ones you are experiencing a serious emotional struggle.

Where can I get help?

Get help from any of these community resources:

- Share Parents of Utah. This organization provides support, services, education, and other resources for grieving parents and siblings at the time of and after the death of a baby. Share Parents of Utah can get involved at any stage of child loss, including miscarriage, stillbirth, or infant death. Call 801-272-5355 or visit share-parents-of-utah.com.
- **Postpartum Support International Utah Chapter**. The Utah Maternal Health Collaborative offers free phone and email support provided by women who have experienced emotional health complications as well as helpful local resources and referrals. Call the crisis line at 801-587-3000 or visit **psiutah.org**.
- **Postpartum Support International.** This agency provides education and resources to women with any of the symptoms listed on <u>page 1</u>. Call 800-944-4773 or visit <u>postpartum.net</u>.
- United Way, Help Me Grow. Call 801-691-5322 and volunteers will link you to community mental health services.
- National Suicide Prevention Hotline. Call 800-273-TALK (8255).
- National Peer Mom Volunteers. Call 800-PPD-MOMS (773-6667).

Questions for my doctor	