

Sodium-Restricted Daily Eating Plan: *2000 milligrams of sodium or less recommended per day*

What is sodium?

Sodium is a mineral that helps your body balance its fluids. Too much sodium in your diet can cause your body to retain fluid. This extra fluid can raise your blood pressure and increase your risk for cardiovascular disease, congestive heart failure, and kidney disease.

Where does all this sodium come from? The major source of sodium in your diet is salt. Some sodium comes from the salt you add to your food from the salt shaker on the table. A lot of salt is also already in processed foods. Reducing these sources of salt can help you significantly lower your sodium intake to under the recommended 2000 milligrams a day. **The goal of this eating plan is to recommend — and help you recognize — those foods that contain less salt.**

How much sodium is okay?

Your healthcare providers are recommending you **eat only 2000 milligrams of sodium or less per day.** The human body really only needs around 500 milligrams of sodium daily (the equivalent of about ¼ teaspoon of salt). Most people, though, get 4000-6000 milligrams daily, almost three teaspoons of salt and around ten times what your body requires! (A single teaspoon of salt alone has about 2300 milligrams of sodium.) Eating less salt will significantly help you reduce the amount of sodium in your diet.



General guidelines

To help you limit the amount of sodium you consume, follow these general guidelines:

- **Don't add salt to food you prepare or eat.** You can use salt substitutes if your physician approves them, but you still need to use them sparingly and not use them during food preparation. You should also avoid "lite salt," which is still a ½ salt mix.
- **Avoid commercially processed foods, which contain large amounts of sodium.**
- **Watch out for medications that contain sodium** — like some sedatives, antacids and alkalizers, headache remedies, or laxatives. If you follow a low-sodium diet, talk to your physician or pharmacist before using these medications.
- **Be extra careful at restaurants** — especially fast food restaurants. Get to know the restaurants in your area and the menu items they offer. Generally, food that's baked or broiled, and made without sauces or gravies, will have less sodium.
- **Avoid sea salt.** It's not healthier than table salt.
- **Avoid soft water whenever possible.** Water softeners may add a lot of sodium to your water. Drink hard water instead. If you drink bottled water check the label for the sodium content.

Dietitian: _____

Phone: _____

Facility: _____

The ABCs of herbs and spices

Instead of reaching for the salt shaker, try some of these sodium-free seasonings to enhance the natural flavors in food:







<i>Allspice</i>	stew, tomatoes, peaches, applesauce, gravies
<i>Basil</i>	fish, lamb, salads, soups, Italian sauces
<i>Cayenne</i>	soups, casseroles, cheese sauces, baked egg dishes, BBQ, poultry and lean meats
<i>Cinnamon</i>	fruits, pork, bread, sweet potatoes, squash, plain yogurt
<i>Cumin</i>	chili, Asian meat dishes, stews, beans
<i>Curry</i>	lean meats, chicken, fish, tomatoes, sauces, rice
<i>Dill</i>	fish, chicken, vegetables, potatoes, salads, pasta
<i>Fennel</i>	lean pork, squash, beets, pasta sauces, seafood dishes
<i>Garlic</i>	lean meats, fish, poultry, soups, salads, vegetables, pasta dishes
<i>Ginger</i>	lean pork, fish, fruits, carrots, pumpkin
<i>Lemon/Lime juice</i>	fish, poultry, salads, vegetables, sauces
<i>Marjoram</i>	shellfish, veal, poultry, soups, stews, eggplant
<i>Mint</i>	lamb, fruit, salads, potatoes, bulgur, beans
<i>Mustard (dry)</i>	lean beef, ham, fish, salads, vegetable dishes
<i>Nutmeg</i>	potatoes, chicken, fish, cauliflower, broccoli, and cabbage dishes
<i>Onion</i>	lean meats, stews, vegetables, salads, soups, legumes
<i>Oregano</i>	sauces, Italian dishes, pizza, salads, vegetables
<i>Paprika</i>	fish, poultry, soups, salads, sauces, baked potatoes, beans
<i>Parsley</i>	cottage cheese, soups, salads, meats, fish, poultry, eggs
<i>Rosemary</i>	chicken, fish, lean pork sauces, stuffing, potatoes, peas, lima beans
<i>Saffron</i>	rice and potato dishes, breads, sauces, soups, vegetables, squash, tomatoes
<i>Savory</i>	salads, lean pork, lean ground meats, soups, vegetables, squash, tomatoes
<i>Sesame</i>	salads, breads, chicken, vegetables, casseroles
<i>Tarragon</i>	sauces, salad dressings, marinades, vegetables, beets, chicken, fish
<i>Thyme</i>	lean meats, poultry, sauces, soups, peas, salads, tomatoes






Herb Hints

- Long cooking times can cause herbs and spices to lose their flavors. Add herbs or spices to cooked foods within 1 hour of serving to get the most flavor.
- Powdered herbs are stronger than dried, and dried are stronger than fresh. As a guide: ¼ teaspoon of powdered herbs gives the flavor of ¾ to 1 teaspoon of dried herbs or 2 teaspoons of fresh herbs.
- You can store dried herbs — covered in a plastic or glass container and in a dry, cool spot out of direct sunlight — for up to a year.
- Wine and herb vinegars make a great marinade for meats and sprinkled on vegetables.
- Lemon, lime, and pineapple juice also make a good marinade that you can sprinkle on vegetables for a unique taste.





FOOD GROUP	FOODS RECOMMENDED	FOODS TO AVOID
<p>Milk and milk products (Limit to 2 cups of milk daily)</p> 	<ul style="list-style-type: none"> • Whole, low fat, or skim <ul style="list-style-type: none"> – cocoa, chocolate, yogurt, or eggnog • Substitute 4 ounces evaporated or condensed milk or 1/3 cup dry milk powder for 8 ounces of milk 	<ul style="list-style-type: none"> • Dutch processed chocolate, buttermilk, malted milk
<p>Breads</p> 	<ul style="list-style-type: none"> • Except those listed in foods to avoid, all whole grain or enriched breads including: <ul style="list-style-type: none"> – homemade biscuits, pancakes, waffles, and muffins, all made without adding salt – unsalted pretzels, low sodium crackers, and low sodium or homemade bread crumbs 	<ul style="list-style-type: none"> • Bread, rolls, or crackers, with visible salted tops • Bread products made using prepared mixes, self-rising flour, or salt • Commercially prepared bread crumbs, croutons, or cracker crumbs
<p>Cereals</p> 	<ul style="list-style-type: none"> • Cooked cereal prepared without salt • Dry cereals: <ul style="list-style-type: none"> – less than 250 milligrams of sodium per serving 	<ul style="list-style-type: none"> • Quick cooking or instant cereals, and all other dry cereals
<p>Fruits</p> 	<ul style="list-style-type: none"> • All fruits and fruit juices 	<ul style="list-style-type: none"> • None
<p>Vegetables</p> 	<ul style="list-style-type: none"> • Fresh, frozen, and low sodium canned vegetables • Low sodium vegetable juices 	<ul style="list-style-type: none"> • Regular canned vegetables and vegetable juices • Sauerkraut, pickled vegetables, or those prepared in brine • Frozen vegetables in a sauce • Regular tomato sauce, puree, and paste

This handout is intended for use during an appointment with a registered dietitian only, and is not meant for use by other clinicians.

FOOD GROUP	FOODS RECOMMENDED	FOODS TO AVOID
<p>Potatoes and starches</p> 	<ul style="list-style-type: none"> • White or sweet potatoes and low salt potato chips • Enriched rice and barley • Noodles, spaghetti, macaroni, and other pastas • Homemade bread stuffing 	<ul style="list-style-type: none"> • Potato casserole mixes and any frozen or dehydrated potatoes • Salted potato chips and salty snacks • Instant rice and pasta mixes • Commercial casserole and stuffing
<p>Meat and meat substitutes</p> 	<ul style="list-style-type: none"> • Any fresh or fresh-frozen beef, lamb, pork, veal, game, and poultry • Any fresh-water or fresh-frozen fish (without breading) and shellfish • Low sodium canned tuna and salmon, or sardines • Unsalted peanut butter, dried beans, and peas • Eggs • Low sodium cheese, cream cheese, ricotta cheese, rinsed and drained cottage cheese 	<ul style="list-style-type: none"> • Smoked, salt-cured, dried, canned, or frozen meat, fish, or poultry • Bacon, ham, cold cuts, hot dogs, sausages, chipped, or corned beef • Kosher-prepared meat • Sardines, anchovies, marinated herring, pickled meats or eggs • Regular canned tuna or salmon • Hard and processed cheese, and cheese spreads • Peanut butter and frozen dinners
<p>Fats</p> 	<ul style="list-style-type: none"> • Unsalted butter or margarine • Unsalted salad dressings and mayonnaise-type salad dressings • Vegetable oil and shortening • Light and heavy cream, sour cream 	<ul style="list-style-type: none"> • Butter, margarine, salad dressings, bacon fat, and salt pork • Snack dips made with cheese, bacon, and buttermilk • Instant soup mixes
<p>Soups</p> 	<ul style="list-style-type: none"> • All unsalted homemade soups and cream soups made with recommended foods and milk allowance (limit milk products to 2 cups per day) • Low sodium broth, bouillon, and consommé • Low sodium canned or dehydrated commercial soups 	<ul style="list-style-type: none"> • Regular bouillon, broth, and consommé • Regular canned or dehydrated commercial soups
<p>Beverages</p> 	<ul style="list-style-type: none"> • All beverages, but: <ul style="list-style-type: none"> – drink milk within the set recommendations 	<ul style="list-style-type: none"> • Commercially softened water, foods or beverages made with softened water • Sports drinks

This handout is intended for use during an appointment with a registered dietitian only, and is not meant for use by other clinicians.

FOOD GROUP	FOODS RECOMMENDED	FOODS TO AVOID
<p>Sweets and desserts</p> 	<ul style="list-style-type: none"> All sweets and desserts except those made from foods that are not recommended With desserts made from milk, the milk needs to be counted as part of the 2 cups daily allowance 	<ul style="list-style-type: none"> Pastries prepared with salted crusts
<p>Other</p> 	<ul style="list-style-type: none"> Pepper, herbs, spices, flavoring, vinegar, lemon or lime juice, and salt-free seasoning mixes The following low sodium condiments: <ul style="list-style-type: none"> catsup, mustard, chili sauce, fresh ground horseradish, Tabasco sauce, and mustard powder Unsalted nuts, seeds, popcorn, and pretzels 	<ul style="list-style-type: none"> Garlic salt, onion salt, celery salt, sea salt, rock salt, and Kosher salt Any other seasoning containing salt and sodium compounds Monosodium glutamate (Accent) Regular: <ul style="list-style-type: none"> catsup, chili sauce, mustard, pickles, relishes, olives, and horseradish Kitchen Bouquet, gravy and sauce mixes BBQ, soy, teriyaki, Worcestershire, and steak sauce Salted nuts, seeds, popcorn, and pretzels All commercially prepared and convenience foods

Nutrition Label Hints

When you see these terms on a food label, here's what they mean:

- Sodium Free** — Less than 5 milligrams of sodium per serving
- Very Low Sodium** — 35 milligrams or less of sodium per serving
- Low Sodium** — 149 milligrams or less of sodium per serving
- Reduced Sodium** — 25% less sodium than the original version of the product
- No Added Salt or Unsalted** — No salt is added during the processing (this does not guarantee the food product is sodium-free)



Intermountain Healthcare complies with applicable federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex. Se proveen servicios de interpretación gratis. Hable con un empleado para solicitarlo. 我們將根據您的需求提供免費的口譯服務。請找尋工作人員協助

This handout is intended for use during an appointment with a registered dietitian only, and is not meant for use by other clinicians.

© 2004-2017 Intermountain Healthcare. All rights reserved. The content presented here is for your information only. It is not a substitute for professional medical advice, and it should not be used to treat a health problem or disease. Please consult your healthcare provider if you have questions or concerns. Patient and Provider Publications RD001 - 01/17