

# Pediatric Dosing Guide: Acetaminophen (Tylenol)

See the packaging that comes with the medicine for complete directions and warnings. Always follow your pediatrician’s directions. Taking too much acetaminophen can cause problems.

Call your pediatrician right away if a child under 3 months has a fever of 100.4° F (38° C) or higher (unless your child received vaccines that day).

Follow these tips to give your child the correct dose of acetaminophen:

- Use your child’s weight and the table below to select the correct dose.
- Do not give acetaminophen for more than 7 days in a row without talking to your pediatrician.
- You can give the doses listed in the table below every 4 hours, if necessary. Give the minimum number of doses necessary to help your child feel better. Do not give more than 5 doses in 24 hours.

- Acetaminophen is found in many over-the-counter and prescription pain medicines. For example, DayQuil contains acetaminophen. Read the labels, and **don’t give your child more than 1 medicine containing acetaminophen at a time** without talking to your child’s doctor.



## Acetaminophen (Tylenol) Dosing Guide

Weight (pounds)	Dose	Children’s liquid syrup (use spoon provided)	Children’s chewable tablets or meltaways	Junior-strength chewable tablets or meltaways
Strength		160 mg / 5 mL	80 mg	160 mg
9–11 lbs	60 mg	1.9 mL		
12–17 lbs	80 mg	2.5 mL		
18–23 lbs	120 mg	3.75 mL		
24–35 lbs	160 mg	5 mL	2 tablets	
36–46 lbs	240 mg	7.5 mL	3 tablets	
47–58 lbs	320 mg	10 mL	4 tablets	2 tablets
59–70 lbs	400 mg	12.5 mL	5 tablets	2½ tablets
70+ lbs	480 mg	15 mL	6 tablets	3 tablets

# Pediatric Dosing Guide: Ibuprofen (Motrin, Advil)

**Do not give ibuprofen to children under 6 months.** See the packaging that comes with the medicine for complete directions and warnings. Always follow your pediatrician's directions. Taking too much ibuprofen can cause problems.

Call your pediatrician immediately if a child under 3 months has a fever of 100.4° F (38° C) or higher.

Follow these tips to give your child the correct dose of ibuprofen:

- Check the bottle to determine which concentration you have. Liquid ibuprofen is available in different concentrations.
- Use your child's weight and the table below to select the correct dose.
- **Do not give ibuprofen for more than 7 days** in a row without talking to your pediatrician.



- **You can give the doses listed in the table below every 6 hours, if necessary.** Give the minimum number of doses needed to help your child feel better. **Do not give more than 4 doses in 24 hours.**

- **Give the least amount of medicine necessary to help your child feel better.** Half of the recommended dose may be effective.

If your child is having surgery within 1 week, **do not give ibuprofen.** It can slightly increase bleeding risk.

## Ibuprofen (Motrin, Advil) Dosing Guide

Weight (pounds)	Dose (up to)	Infant drops (use dropper provided)	Children's liquid (use syringe or spoon provided)	Junior-strength chewables	Junior-strength swallow tablets	Adult regular-strength tablets
<b>Concentration</b>		<b>50 mg / 1.25 mL</b>	<b>100 mg / 5 mL</b>	<b>100 mg</b>	<b>100 mg</b>	<b>200 mg</b>
11–16 lbs	<b>50 mg</b>	1.25 mL	2.5 mL			
17–21 lbs	<b>75 mg</b>	1.9 mL	3.75 mL			
22–32 lbs	<b>100 mg</b>		5 mL	1 tablet		
33–43 lbs	<b>150 mg</b>		7.5 mL	1½ tablets		
44–54 lbs	<b>200 mg</b>		10 mL	2 tablets	2 tablets	
55–65 lbs	<b>250 mg</b>		12.5 mL	2½ tablets	2½ tablets	1 tablet
66–87 lbs	<b>300 mg</b>		15 mL	3 tablets	3 tablets	1 to 1½ tablets
88+ lbs	<b>350 mg</b>		17.5 mL	3½ tablets	3½ tablets	1½ to 2 tablets

Intermountain Health complies with applicable federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex. Se proveen servicios de interpretación gratis. Hable con un empleado para solicitarlo. 我們將根據您的需求提供免費的口譯服務。請找尋工作人員協助。

© 2015–2024 Intermountain Health. All rights reserved. The content presented here is for your information only. It is not a substitute for professional medical advice, and it should not be used to diagnose or treat a health problem or disease. Please consult your healthcare provider if you have any questions or concerns. More health information is available at [intermountainhealth.org](http://intermountainhealth.org). CLD003 - 02/24 (Last reviewed - 02/24) Also available in Spanish.