


# Suicide Prevention – CLINIC

Reference  
Link

## SCREENING and ACTIONS

(+) on PHQ-9 (q. 9) or clinical suspicion of suicidal ideation or behaviors





# Suicide Prevention – CLINIC

[Reference Link](#)

## ADDITIONAL SCREENING TOOLS

Columbia Suicide Severity Rating Scale (C-SSRS)		
C-SSRS version	Description	Who administers?
<u><i>Adult/Adolescent (≥12 years) Quick Screen</i></u>	Used in primary care and EDs to quickly screen patients for suicidal thoughts and behaviors; 3–6 questions (depending on patient responses). <i>Assesses past month.</i>	Any clinician can administer this screen.  At Intermountain, the Quick Screen is used by: <ul style="list-style-type: none"> <li>• Primary care providers (PCPs)</li> <li>• Emergency department (ED) triage nurses</li> <li>• Clinicians (nurses, MDs, etc.) on BH units and inpatient units</li> </ul>
<u><i>Pediatric (≤11 years) Quick Screen</i></u>		
<u><i>Adult/Adolescent (≥12 years) Lifetime/Recent Assessment</i></u>	Used for full assessment during initial visit; 2-page assessment (number of questions varies based on patient responses). <i>Assesses lifetime and past month.</i>	<ul style="list-style-type: none"> <li>• PCPs treating patients at low risk</li> <li>• Any MD</li> <li>• Mental health specialists</li> <li>• Trained MHI team members (care managers, health advocates, etc.)</li> <li>• ED crisis workers</li> <li>• Other trained clinicians</li> </ul>
<u><i>Pediatric (≤11 years) Lifetime/Recent Assessment</i></u>		
<u><i>C-SSRS — Adult/Adolescent (≥12 years) Since Last Visit Version</i></u>	Used to assess patient at follow-up visits; same questions as the Lifetime/Recent versions. <i>Assesses since last visit or since last assessment.</i>	
<u><i>Pediatric (≤11 years) Since Last Visit Version</i></u>		
<u><i>Suicide Prevention — Risk Assessment Tool</i></u>	Used to assess the patient's level of risk during initial visit and over time; 1-page list of suicide risk and protective factors. <i>Assesses current risk.</i>	