

# Heart Failure: Tracking your fluids

When you have heart failure, you need to limit the sodium (salt) and fluids that you drink and eat each day. This is especially true for people who retain fluid and need to take water pills. This sheet will show you simple ways to track your fluids each day.

## Why do I need to track my fluids?

Limiting fluids can ease swelling (edema) and lighten the workload on your heart. Generally, limit the amount of fluids that you drink or eat to around 2 liters (2,000 milliliters) per day. (Check with your healthcare provider to be sure.)

Use these tips to help limit your fluids:

- **Sip a small bottle of water** throughout the day (record it in your diary as part of a snack).
- Fight thirst by using lip salve/gloss, sucking on ice chips, eating frozen grapes or hard candies, chewing sugar-free gum, or using mouth rinses or swabs.

Use the table on the right to learn how to convert ounces to milliliters.

#### Conversion Table

Common amounts	Ounces	mL
2 tablespoons	<b>1</b> oz	<b>30</b> mL
1 small juice glass	<b>4</b> oz	<b>120</b> mL
1 cup	<b>8</b> oz	<b>240</b> mL
1 cup of ice chips (Ice chips melt to half the fluid)	<b>4</b> oz	<b>120</b> mL
1 can of soda	<b>12</b> oz	<b>360</b> mL
TARGET: Daily amount less than	<b>64</b> oz	<b>2,000</b> mL

Track your fluids using the same units of measure each time. This will make it easy to add up your daily amount. The table above shows ounce (oz) and milliliter (mL) equivalents for common amounts. Use the formula "1 oz = 30 mL" to figure the mL for any amount in ounces.

## Know how much you drink

Take a look at the cup you drink from most often. How much does it hold? Fill it up to the level that you usually use. Pour that amount into a measuring cup. Then use the conversion formula (1 ounce = 30 mL) to figure how much you are drinking and write it here. My cup holds \_\_\_\_\_\_ ounces (or \_\_\_\_\_ mL).

How many times can you fill your cup in a day and still stay under 64 ounces (or 2,000 mL)?

See the other side of this sheet for an easy way to track your 2 liters of fluid.









### Tracking your 2 liters of fluids


Follow the steps below to help you keep track of how many fluids you take in each day.

1 Litre


**Step 1**: Fill a 2-liter bottle with 64 ounces (1.9 liters) of water and set it next to your kitchen sink. Place a measuring cup nearby.



**Step 2**: Each time you eat or drink, figure out how much fluid you had. This can be from something you drank or from food that you ate (see examples below).









Fluids	Ounces	mL	Fluids	Ounces	mL
Watermelon, 1 cup	4.7 oz	140 mL	Chicken noodle soup (low sodium) 1 cup	7.5 oz	222 mL
Strawberries, 1 cup,	4.6 oz	138 mL	Spaghetti sauce (low sodium), ½ cup	4 oz	117 mL
Romaine lettuce, 1 cup (shredded)	1.5 oz	45 mL	Celery (raw), 1 cup	3.2 oz	96 mL
Tomato, 3 medium slices	2 oz	57 mL	Cottage cheese, 1 cup	4 oz	117 mL
Cucumber slices, 1 cup	3.3 oz	100 mL	Jello, ½ cup	3.8 oz	113 mL
Ice cream, ½ cup	1.5 oz	45 mL	Popsicle, 1	1.75 oz	52 mL
Yogurt, 6 oz container	4.3 oz	128 mL	Pudding, ½ cup	3.7 oz	108 mL
My foods:			My foods:		
My foods:			My foods:		

Intermountain Healthcare complies with applicable federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex. Se proveen servicios de interpretación gratis. Hable con un empleado para solicitarlo. 我們將根據您的需求提供免費的口譯服務。請找尋工作人員協助。