


Suicide Prevention – ED

Reference
Link

SCREENING and ACTIONS

Patient presents with mental health concern or clinical suspicion of suicidal ideation present



Suicide Prevention – ED

Reference
Link

ADDITIONAL SCREENING TOOLS

Columbia Suicide Severity Rating Scale (C-SSRS)		
C-SSRS version	Description	Who administers?
<u>Adult/Adolescent (≥12 years) Quick Screen</u>	Used in primary care and EDs to quickly screen patients for suicidal thoughts and behaviors; 3–6 questions (depending on patient responses). <i>Assesses past month.</i>	Any clinician can administer this screen. At Intermountain, the Quick Screen is used by: <ul style="list-style-type: none"> • Primary care providers (PCPs) • Emergency department (ED) triage nurses • Clinicians (nurses, MDs, etc.) on BH units and inpatient units
<u>Pediatric (≤11 years) Quick Screen</u>		
<u>Adult/Adolescent (≥12 years) Lifetime/Recent Assessment</u>	Used for full assessment during initial visit; 2-page assessment (number of questions varies based on patient responses). <i>Assesses lifetime and past month.</i>	<ul style="list-style-type: none"> • PCPs treating patients at low risk • Any MD • Mental health specialists • Trained MHI team members (care managers, health advocates, etc.) • ED crisis workers • Other trained clinicians
<u>Pediatric (≤11 years) Lifetime/Recent Assessment</u>		
<u>C-SSRS — Adult/Adolescent (≥12 years) Since Last Visit Version</u>		
<u>Pediatric (≤11 years) Since Last Visit Version</u>	Used to assess patient at follow-up visits; same questions as the Lifetime/Recent versions. <i>Assesses since last visit or since last assessment.</i>	
<u>Suicide Prevention — Risk Assessment Tool</u>	Used to assess the patient's level of risk during initial visit and over time; 1-page list of suicide risk and protective factors. <i>Assesses current risk.</i>	